arXiv:1612.09156v1 [math.AG] 29 Dec 2016

MATHER-JACOBIAN MULTIPLIER IDEALS ON TORIC VARIETIES

HOWARD M THOMPSON

Abstract. This paper provides a formula for the Mather-Jacobian multiplier ideals of torus invariant ideals on (not necessarily normal) toric varieties that generalizes HowaldâĂŹs formula for the multiplier ideal of monomial ideals in a polynomial ring.

1. Introduction

In Section 2, we recall the definition of Nash blowup and the relationship between Nash blowups and Jacobian ideals. In Section 3 on the next page, we recall the definition of logarithmic Jacobian ideals and the relationships between Nash blowups, logarithmic Jacobian ideals, and Jacobian ideals. In Section 4 on page 3, we recall the definition of Mather-Jacobian multiplier ideals and present a formula for the Mather-Jacobian multiplier ideals of torus invariant ideals on (not necessarily normal) toric varieties.

2. Nash Blowup and the Jacobian

In this section we loosely follow de Fernex & Docampo [2, Section 2].

Let X be a separated reduced scheme of finite type over an algebraically closed field $\Bbbk$ of characteristic zero. The *Jacobian ideal sheaf* of X, $\mathcal{J}ac_X \subseteq \mathcal{O}_X$, is the smallest non-zero Fitting ideal of the cotangent sheaf Ω_X. If X is of pure dimension d, then $\mathcal{J}ac_X = \mathrm{Fitt}^d(\Omega_X)$ is the d-th Fitting ideal of Ω_X. If Y is a smooth scheme and $f : Y \to X$ is a morphism of schemes, the Jacobian ideal of f is the 0-th Fitting ideal $\mathcal{J}ac_f = \mathrm{Fitt}^0(\Omega_{Y/X})$ of $\Omega_{Y/X}$. If Y is equidimensional of dimension d, then Ω^d_Y is an invertible sheaf and the image of the map $f^*\Omega^d_X \to \Omega^d_Y$ is $\mathcal{J}ac_f \cdot \Omega^d_Y$.

Let X be a separated reduced scheme of finite type over $\Bbbk$ of pure dimesion d. The *Nash blowup* of X is the birational map $\pi : \widehat{X} \to X$ satisfying the universal property: if $\mu : Y \to X$ is a proper birational map such that $\mu^*\Omega^1_X/(torsion)$ is a locally free $\mathcal{O}_Y$-module, then μ factors uniquely through π. Following Ein, Ishii & Mustaţă [3, Remark 2.3] and noticing that X need not be a variety to apply Lipman [6, Lemma 1], we see every log resolution of $\mathcal{J}ac_X$ factors through the Nash blowup of X. In fact, by Oneto & Zatini [7, Corollary 5.2],

Date: December 30, 2016.
2010 *Mathematics Subject Classification.* Primary: 14F18; Secondary: 14M25.
Key words and phrases. Multiplier ideals, toric varieties.

if $\mathcal{J}ac_Y|_X$ is the restriction to X of the Jacobian of a locally complete intersection Y of dimension d containing X, then the Nash blowup is the blowup of $\mathcal{J}ac_Y|_X$.

3. The Jacobian and the Logarithmic Jacobian

From now on, let S be an affine semigroup (finitely generated torsion-free cancellative monoid) and let $M = \mathbb{Z}\mathsf{S} \cong \mathbb{Z}^d$ be its quotient group. So, $X = \operatorname{Spec}\Bbbk[\mathsf{S}]$ is a (not necessarily normal) affine toric variety. We write $\mathbf{m}$ for an element of $M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$ and we write $\chi^{\mathbf{m}}$ for the element of $\Bbbk[M]$ corresponding to $\mathbf{m} \in M$. We have a $\Bbbk[\mathsf{S}]$-module homomorphism

$$\varphi : \Omega^1_{\Bbbk[\mathsf{S}]/\Bbbk} \to \Bbbk[\mathsf{S}] \otimes_{\mathbb{Z}} M; \quad \varphi(d\chi^{\mathbf{m}}) = \chi^{\mathbf{m}} \otimes \mathbf{m}.$$

Definition 1. The *logarithmic Jacobian ideal* $\mathcal{J}ac_X^{log}$ of X is the ideal such that the image of $\wedge^d\varphi$ is $\mathcal{J}ac_X^{log} \otimes_{\mathbb{Z}} \wedge^d M$. This ideal is generated by elements $\chi^{\mathbf{m}_1+\mathbf{m}_2+\cdots+\mathbf{m}_d}$ such that $\mathbf{m}_1 \wedge \mathbf{m}_2 \wedge \cdots \wedge \mathbf{m}_d \neq \mathbf{0}$. Evidently, it suffices to take the $\mathbf{m}_i$ from some fixed generating set for S.

According to González Pérez & Teissier [4, Proposition 60], the Nash blowup $\pi : \widehat{X} \to X$ of X is the blowup of $\mathcal{J}ac_X^{log}$. Examining the proof of this proposition provides a convenient formula for the Jacobian of X. Fix a surjective monoid homomorphism $\phi : \mathbb{N}^r \to \mathsf{S}$. This surjection induces a short exact sequence of Abelian groups

$$0 \longrightarrow \mathcal{L} \longrightarrow \mathbb{Z}^r \xrightarrow{\phi^{gp}} M \longrightarrow 0\ .$$

For each element $\mathbf{u} \in \mathbb{Z}^r$, write $\mathbf{u} = \mathbf{u}^+ - \mathbf{u}^-$ where $\mathbf{u}^+$ and $\mathbf{u}^-$ are elements of $\mathbb{N}^r$ with disjoint support. The monoid homomorphism ϕ induces a $\Bbbk$-algebra homomorphism $\phi^{alg} : \Bbbk[x_1, x_2, \ldots, x_r] \to \Bbbk[\mathsf{S}]$ with kernel $I_{\mathcal{L}} = \left(\mathbf{x}^{\mathbf{u}^+} - \mathbf{x}^{\mathbf{u}^-} \mid \mathbf{u} \in \mathcal{L}\right)$. Set $c = \operatorname{rank}\mathcal{L} = r - d$, let $\mathbf{e}_1, \mathbf{e}_2, \ldots \mathbf{e}_r$ be the standard basis in $\mathbb{N}^r$, let $\mathbf{1} = \sum_{i=1}^r \mathbf{e}_i$ be the all ones vector, and fix a generating set $\left\{f_1 = \mathbf{x}^{\mathbf{u}_1^+} - \mathbf{x}^{\mathbf{u}_1^-}, f_2 = \mathbf{x}^{\mathbf{u}_2^+} - \mathbf{x}^{\mathbf{u}_2^-}, \ldots, f_n = \mathbf{x}^{\mathbf{u}_n^+} - \mathbf{x}^{\mathbf{u}_n^-}\right\}$ for $I_{\mathcal{L}}$.

Let J be the $r \times n$ matrix $\left[\frac{\partial f_j}{\partial x_i}\right]$, let U be the $r \times n$ matrix with columns $\mathbf{u}_1, \mathbf{u}_2, \ldots, \mathbf{u}_n$, let $K = \{k_1, k_2, \ldots, k_c\} \subseteq \{1, 2, \ldots, r\}$, and let $L = \{j_1, j_2, \ldots, j_c\} \subseteq \{1, 2, \ldots, n\}$. González Pérez & Teissier prove following relationship between the $c \times c$ minors J_{KL} and U_{KL}.

$$\mathbf{x}^{\mathbf{e}_{k_1}+\mathbf{e}_{k_2}+\cdots\mathbf{e}_{k_c}} \cdot J_{KL} \equiv \mathbf{x}^{\mathbf{u}_{j_1}^+ + \mathbf{u}_{j_2}^+ + \cdots + \mathbf{u}_{j_c}^+} U_{KL} \mod I_{\mathcal{L}}. \tag{1}$$

Now, fix $1 \leq j_1 < j_2 < \cdots < j_c \leq n$ such that $\mathbf{0} \neq \mathbf{u}_{j_1} \wedge \mathbf{u}_{j_2} \wedge \cdots \wedge \mathbf{u}_{j_c} \in \wedge^c\mathcal{L}$. Let $I = \left(\mathbf{x}^{\mathbf{u}_{j_1}^+} - \mathbf{x}^{\mathbf{u}_{j_1}^-}, \mathbf{x}^{\mathbf{u}_{j_2}^+} - \mathbf{x}^{\mathbf{u}_{j_2}^-}, \ldots, \mathbf{x}^{\mathbf{u}_{j_c}^+} - \mathbf{x}^{\mathbf{u}_{j_c}^-}\right) \subseteq \Bbbk[\mathbf{x}]$ Then, $Y = \operatorname{Spec}(\Bbbk[\mathbf{x}]/I)$ is a reduced complete intersection of dimension d containing X. After multiplying both sides of the congruence Eq. (1) by $\mathbf{x}^{\mathbf{1}-(\mathbf{e}_{k_1}+\mathbf{e}_{k_2}+\cdots\mathbf{e}_{k_c})}$, varying K, and considering the restriction to X, one obtains

$$\chi^{\phi^{gp}(\mathbf{1})} \cdot \mathcal{J}ac_{Y|X} = \chi^{\phi^{gp}(\mathbf{u}_{j_1}^+ + \mathbf{u}_{j_2}^+ + \cdots + \mathbf{u}_{j_c}^+)} \cdot \mathcal{J}ac_X^{log}.$$

Lemma 2. *$\mathcal{J}ac_X$ is generated by*

$$\left\{ \chi^{\phi^{gp}(\mathbf{e}_{i_1}+\mathbf{e}_{i_2}+\cdots+\mathbf{e}_{i_d}+\mathbf{u}_{j_1}^+ +\mathbf{u}_{j_2}^+ +\cdots+\mathbf{u}_{j_c}^+ -\mathbf{1})} \;\middle|\; \begin{array}{c} \mathbf{0} \neq \phi^{gp}(\mathbf{e}_{i_1}) \wedge \phi^{gp}(\mathbf{e}_{i_2}) \wedge \cdots \wedge \phi^{gp}(\mathbf{e}_{i_d}) \in \wedge^d M; \\ \mathbf{0} \neq \mathbf{u}_{j_1} \wedge \mathbf{u}_{j_2} \wedge \cdots \wedge \mathbf{u}_{j_c} \in \wedge^c \mathcal{L} \end{array} \right\}.$$

Proof. Continuing as above, allow L to vary as well. We deduce

$$\chi^{\phi^{gp}(\mathbf{1})} \cdot \mathcal{J}ac_X = \mathcal{J}ac_X^{log} \cdot \left(\chi^{\phi^{gp}(\mathbf{u}_{j_1}^+ +\mathbf{u}_{j_2}^+ +\cdots+\mathbf{u}_{j_c}^+)} \;\middle|\; \mathbf{0} \neq \mathbf{u}_{j_1} \wedge \mathbf{u}_{j_2} \wedge \cdots \wedge \mathbf{u}_{j_c} \in \wedge^c \mathcal{L} \right).$$

□

4. Mather-Jacobian Multiplier Ideals

Definition 3. Let $\mathfrak{a} \subseteq \mathcal{O}_X$ be a nonzero ideal sheaf on a variety X, let $\mu : Y \to X$ be a log resolution of $\mathfrak{a} \cdot \mathcal{J}ac_X$, let $\mathcal{O}_Y\left(-\widehat{K}_{Y/X}\right) = \mathcal{J}ac_\mu$, let $\mathcal{O}_Y(-J_{Y/X}) = \mathcal{J}ac_X \cdot \mathcal{O}_Y$, let $\mathcal{O}_Y(-Z_{Y/X}) = \mathfrak{a} \cdot \mathcal{O}_Y$, and let $\lambda \in \mathbb{R}_{\geq 0}$. The *Mather-Jacobian multiplier ideal* of $\mathfrak{a}$ with exponent λ is

$$\widehat{\mathcal{J}}(X, \mathfrak{a}^\lambda) = \mu_* \mathcal{O}_Y \left(\widehat{K}_{Y/X} - J_{Y/X} - \lfloor \lambda Z_{Y/X} \rfloor \right).$$

Now, we are ready to develop a formula for the Mather-Jacobian multiplier ideals of torus invariant ideals on (not necessarily normal) toric varieties. it suffice to consider the affine case. Let $X = \operatorname{Spec} \Bbbk[\mathsf{S}]$ be a (not necessarily normal) affine toric variety, let $\mathfrak{a} \subseteq \Bbbk[\mathsf{S}]$ be a nonzero monomial ideal, let $\mu : X_\Sigma \to X$ be a toric log resolution of $\mathfrak{a} \cdot \mathcal{J}ac_X$.

Lemma 4. *If $\mathfrak{a} \subseteq \mathcal{O}_X$ is a nonzero invariant ideal sheaf on a toric variety X and $\mu : X_\Sigma \to X$ is a toric log resolution of $\mathfrak{a} \cdot \mathcal{J}ac_X$, then*

$$\mathcal{J}ac_\mu \cdot \mathcal{J}ac_{X_\Sigma}^{log} = \mu^* \mathcal{J}ac_X^{log} \text{ and } \mathcal{O}_{X_\Sigma}\left(\widehat{K}_{X_\Sigma/X}\right) = \mathcal{H}om_{\mathcal{O}_{X_\Sigma}}\left(\mu^* \mathcal{J}ac_X^{log}, \mathcal{J}ac_{X_\Sigma}^{log}\right).$$

Proof. Consider the map

$$\begin{array}{ccc} \mu^*\Omega_X^d & \longrightarrow & \Omega_{X_\Sigma}^d \\ \| & & \| \\ \mu^*(\mathcal{J}ac_X^{log} \otimes_{\mathbb{Z}} \wedge^d M) & \longrightarrow & \mathcal{J}ac_{X_\Sigma}^{log} \otimes_{\mathbb{Z}} \wedge^d M \end{array}$$

with image $\mathcal{J}ac_\mu \cdot \Omega_{X_\Sigma}^d = \mathcal{J}ac_\mu \cdot \mathcal{J}ac_{X_\Sigma}^{log} \otimes_{\mathbb{Z}} \wedge^d M$. Since $\mu^* \mathcal{J}ac_X^{log}$, $\mathcal{J}ac_{X_\Sigma}^{log}$ and $\mathcal{J}ac_\mu$ are line bundles on X_Σ, the result follows. □

Recall the notation of Section 3, let $\mathfrak{a} \subseteq \Bbbk[\mathsf{S}]$ be a nonzero monomial ideal, and consider the following subsets of $M_{\mathbb{R}}$.

$$\begin{aligned}
\sigma^\vee &= \operatorname{Conv}(\mathsf{S}) \\
\mathsf{J}^{log} &= \{\phi^{gp}(\mathbf{e}_{i_1} + \mathbf{e}_{i_2} + \cdots + \mathbf{e}_{i_d}) \mid \phi^{gp}(\mathbf{e}_{i_1}) \wedge \phi^{gp}(\mathbf{e}_{i_2}) \wedge \cdots \wedge \phi^{gp}(\mathbf{e}_{i_d}) \neq \mathbf{0}\} \\
\operatorname{Newt}\left(\mathcal{J}ac_X^{log}\right) &= \operatorname{Conv}(\mathsf{J}^{log} + \mathsf{S}) \\
\mathsf{J}' &= \{\phi^{gp}(\mathbf{u}_{j_1}^+ + \mathbf{u}_{j_2}^+ + \cdots + \mathbf{u}_{j_c}^+ - \mathbf{1}) \mid \mathbf{u}_{j_1} \wedge \mathbf{u}_{j_2} \wedge \cdots \wedge \mathbf{u}_{j_c} \neq \mathbf{0}\} \\
\mathsf{J} &= \mathsf{J}^{log} + \mathsf{J}' \\
\operatorname{Newt}\left(\mathcal{J}ac_X\right) &= \operatorname{Conv}(\mathsf{J} + \mathsf{S}) \\
P &= \operatorname{Newt}(\mathfrak{a}) = \operatorname{Conv}(\{\mathbf{m} \mid \chi^{\mathbf{m}} \in \mathfrak{a}\}) \\
Q &= \operatorname{Conv}(\mathsf{J}' + \mathsf{S}) = \left\{\mathbf{m} \mid \mathbf{m} + \operatorname{Newt}\left(\mathcal{J}ac_X^{log}\right) \subseteq \operatorname{Newt}\left(\mathcal{J}ac_X\right)\right\}
\end{aligned}$$

Let $\mathsf{S}_\sigma = \sigma^\vee \cap M$. Thus, $X_\sigma = \operatorname{Spec} \Bbbk[\mathsf{S}_\sigma]$ is a normal toric variety and the canonical map $X_\sigma \to X$ induced by the inclusion $\mathsf{S} \subseteq \mathsf{S}_\sigma$ is the normalization map. We say a map $\mu : X_\Sigma \to X$ from a normal toric variety is *toric* if if factors as a toric map $X_\Sigma \to X_\sigma$ followed by the canonical map $X_\sigma \to X$.

Theorem 5. *If $\mathfrak{a} \subseteq \mathcal{O}_X$ is a nonzero invariant ideal sheaf on a toric variety X and $\mu : X_\Sigma \to X$ is a toric log resolution of $\mathfrak{a} \cdot \mathcal{J}ac_X$, then*

$$\chi^{\mathbf{m}} \in \widehat{\mathcal{J}}(X, \mathfrak{a}^\lambda) \Leftrightarrow \mathbf{m} \in \operatorname{int}(Q + \lambda P),$$

where $\operatorname{int}(Q + \lambda P)$ is the interior of the polyhedron $Q + \lambda P$.

Proof. We will mimic the proof of Howald's Theorem [5] as it appears in Cox, Little & Schenck [1, Theorem 11.3.12]. Let $\Sigma(1)$ be the set of rays of the fan Σ,let $\mathbf{n}_\rho$ be the primitive vector on the ray ρ for each $\rho \in \Sigma(1)$, let D_ρ be the prime divisor on X_Σ for each $\rho \in \Sigma(1)$, and let $Z_{Y/X} = \sum_{\rho \in \Sigma(1)} a_\rho D_\rho$. Exploiting the facts that $\mathcal{J}ac_{X_\Sigma}^{log} = \sum_{\rho \in \Sigma(1)} D_\rho$, $\mu^* \mathcal{J}ac_X^{log} = \sum_{\rho \in \Sigma(1)} \min_{\mathbf{m} \in \mathsf{J}^{log}} \langle \mathbf{m}, \mathbf{n}_\rho \rangle D_\rho$, and $J_{Y/X} = \sum_{\rho \in \Sigma(1)} \min_{\mathbf{m} \in \mathsf{J}} \langle \mathbf{m}, \mathbf{n}_\rho \rangle D_\rho$ along with Lemma 4 on the preceding page, we see

$$\widehat{K}_{Y/X} - J_{Y/X} - \lfloor \lambda Z_{Y/X} \rfloor = \sum_{\rho \in \Sigma(1)} \left(-1 + \min_{\mathbf{m} \in \mathsf{J}^{log}} \langle \mathbf{m}, \mathbf{n}_\rho \rangle - \min_{\mathbf{m} \in \mathsf{J}} \langle \mathbf{m}, \mathbf{n}_\rho \rangle - \lfloor \lambda a_\rho \rfloor \right) D_\rho.$$

Thus,

$$\begin{aligned}
\chi^{\mathbf{m}} \in \widehat{\mathcal{J}}(X, \mathfrak{a}^\lambda) &\Leftrightarrow \langle \mathbf{m}, \mathbf{n}_\rho \rangle \geq 1 - \min_{\mathbf{m} \in \mathsf{J}^{log}} \langle \mathbf{m}, \mathbf{n}_\rho \rangle + \min_{\mathbf{m} \in \mathsf{J}} \langle \mathbf{m}, \mathbf{n}_\rho \rangle + \lfloor \lambda a_\rho \rfloor \qquad \text{for all } \rho \in \Sigma(1) \\
&\Leftrightarrow \langle \mathbf{m}, \mathbf{n}_\rho \rangle > - \min_{\mathbf{m} \in \mathsf{J}^{log}} \langle \mathbf{m}, \mathbf{n}_\rho \rangle + \min_{\mathbf{m} \in \mathsf{J}} \langle \mathbf{m}, \mathbf{n}_\rho \rangle + \lambda a_\rho \qquad \text{for all } \rho \in \Sigma(1) \\
&\Leftrightarrow \langle \mathbf{m}, \mathbf{n}_\rho \rangle > \min_{\mathbf{m} \in \mathsf{J}'} \langle \mathbf{m}, \mathbf{n}_\rho \rangle + \lambda a_\rho \qquad \text{for all } \rho \in \Sigma(1)
\end{aligned}$$

Since the resolution $\mu : X_\Sigma \to X$ principalizes both $\mathcal{J}ac_X^{log}$ and $\mathcal{J}ac_X$, it also principalizes the fractional ideal $(\chi^{\mathbf{m}} \mid \mathbf{m} \in \mathsf{J}')$. Therefore, every facet of $Q + \lambda P$ is supported on a hyperplane of the form $\{\mathbf{m} \mid \langle \mathbf{m}, \mathbf{n}_\rho \rangle = b_\rho\}$ for some $\rho \in \Sigma(1)$ and $b_\rho \in \mathbb{R}$. We conclude

$$\chi^{\mathbf{m}} \in \widehat{\mathcal{J}}(X, \mathfrak{a}^\lambda) \Leftrightarrow \mathbf{m} \in \operatorname{int}(Q + \lambda P).$$

□

References

[1] D. A. Cox, J. B. Little, and H. K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011. MR2810322 ↑4

[2] T. de Fernex and R. Docampo, *Jacobian discrepancies and rational singularities*, J. Eur. Math. Soc. (JEMS) **16** (2014), no. 1, 165–199, DOI 10.4171/JEMS/430. MR3141731 ↑1

[3] L. Ein, S. Ishii, and M. Mustaţă, *Multiplier ideals via Mather discrepancy* (2011), 17 pp., arXiv:1107.2192 [math.AG]. ↑1

[4] P. D. González Pérez and B. Teissier, *Toric geometry and the Semple-Nash modification*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM **108** (2014), no. 1, 1–48, DOI 10.1007/s13398-012-0096-0. MR3183106 ↑2

[5] J. A. Howald, *Multiplier ideals of monomial ideals*, Trans. Amer. Math. Soc. **353** (2001), no. 7, 2665–2671 (electronic), DOI 10.1090/S0002-9947-01-02720-9. MR1828466 ↑4

[6] J. Lipman, *On the Jacobian ideal of the module of differentials*, Proc. Amer. Math. Soc. **21** (1969), 422–426. MR0237511 ↑1

[7] A. Oneto and E. Zatini, *Remarks on Nash blowing-up*, Rend. Sem. Mat. Univ. Politec. Torino **49** (1991), no. 1, 71–82 (1993). Commutative algebra and algebraic geometry, II (Italian) (Turin, 1990). MR1218672 ↑1

Mathematics Department, 402 Murchie Science Building, 303 East Kearsley Street, Flint, MI 48502-1950

E-mail address: hmthomps@umflint.edu

URL: http://homepages.umflint.edu/~hmthomps/